

Inter-Relation between Type 2 Diabetes Mellitus and Hypertension in Asia

Sarathlal S, DNB, Aleena Thomas Cheeran, DNB, Jayakrishnan K S, MD, Hariharan A, DFM, Jamila Hameed *, MD

Department of Medicine, Karuna Medical College, Vilayodi, Kerala, 678103, India.

*Corresponding Author: Jamila Hameed, MD; hameedjamila78@gmail.com

Received: 09 January 2025;

Revised: 17 February 2025;

Accepted: 14 March 2025;

Published: 20 March 2025

Abstract

Type 2 Diabetes Mellitus (T2DM) and hypertension (HTN) are most commonly inter-related. Diabetic patients are at a high risk of developing uncontrolled blood pressure leading to other complications like cardio vascular disease (CVD) and chronic kidney disease (CKD). The risk factors are ageing, obesity, dyslipidaemia, sedentary lifestyle, urbanization, and smoking. A keen comprehension of these interconnected factors will assist in prevention and management. The primary objective of the study is to evaluate the prevalence of HTN in diabetic patients in Asia to establish the inter-relationship to make the general public aware of the condition. The question that we aimed to answer was: "What is the relationship between T2DM and HTN and other variables causing morbidity?". A total of 27103 studies were retrieved from PubMed, Google Scholar and Embase in the period 2014 to 2024 out of which finally 12 studies were considered. HTN prevalence was found to be the highest among T2DM patients in Kuwait and China (87.16% each). Saudi (23%) reported the lowest prevalence followed by Iran (36.1%). The majority of studies reported higher prevalence in females in comparison to males. Factors such as obesity, dyslipidaemia, and low physical activity proved to be the key drivers in the development and progression of diabetes along with urbanization. HTN is a global burden and there has been a strong correlation between T2DM and HTN. Awareness, timely diagnosis and lifestyle modification can prove to be extremely beneficial in appropriate management of the condition and avert mortality.

Keywords: Hypertension, Diabetes Mellitus, Association, Asia.

Introduction

Diabetes is progressively increasing so there is an increase in prevalence of hypertension (HTN). This is the ultimate cause of premature death in developing countries like India. The burden of hypertension and diabetes mellitus (DM) is escalating throughout the globe, especially in Asian countries. The prevalence of hypertension is higher in diabetic patients when compared to non-diabetic. The premature death among the diabetic hypertensive patients is due to cardiovascular disorders amounting to 17 million per year which is one-third of the total deaths in the world. It was estimated in a study that the global estimate for T2DM patients would be around 300 million by 2025 (Zimmet P, 2003). Another study reported the global estimate to be around 366 million by 2030 (Wild S *et al*, 2004). As such over half a million people are currently suffering from diabetes mellitus in Asia. The tragic story is that India is the diabetic capital.

The development of hypertension among diabetic patients includes multiple variables like age, gender, marital status, family history, lack of exercise, inadequate diet, smoking, and BMI. The renal disorders and cardiovascular complications occur in T2DM with HTN (Wang Z *et al*, 2020). About 20-60% diabetic hypertensive patients are obese (Cheung B M, 2010). T2DM with HTN leads to microalbuminuria, retinopathy and it is vital to understand the pathophysiology in detail (Parving H H *et al*, 1988). Hypertension

is a prevalent disease in elderly patients. International Diabetes Federation reports expenditure for diabetic population will reach 1.054 billion USD by 2045 bringing down the global economy (Sun H *et al*, 2022). The T2DM prevalence is estimated to be higher in developing countries due to urbanization by 13.9% which may also be attributed to population ageing. The top ten countries who have Type 2 diabetes mellitus in 2045 were noted reporting that China and India will top the list among them reaching 174 million by 2045 due to urbanization (Cohen B, 2006). The slum residents will also show an increase in prevalence (Devi B N *et al*, 2014). Hypertension prevalence in type 2 diabetes mellitus is mainly attributed to hyperinsulinemia resulting in retention of sodium and water which leads to sympathetic over activity and alteration in the cell membrane thereby altering the cation transport (Sechi LA, Bartoli E, 1997). In addition to this, there is increased angiotensin II and aldosterone activity. Above all oxidative stress is the cornerstone in etiopathology due to tumour necrosis factor and interleukin 6 and plasmin activator-inhibitor. Finally, these changes influence insulin resulting in insulin resistance which is a hallmark in prevalence of hypertension. In our study, we aimed to establish the interrelation between T2DM and HTN and other variables whose presence trigger morbidity.

Materials And Methods

A comprehensive literature review study was done for a period of 11 years from 2014 to 2024. The study was undertaken by two authors (S.S. and A.T.C.) using the keywords “Inter-relation”, “Type 2 Diabetes”, “Hypertension” and “Asia” from PubMed, Google Scholar and Embase. A total of 27103 studies were retrieved and 12 studies were considered finally for the systematic review and meta analyses (**Figure 1**) (Moher D *et al*, 2009).

The inclusion criteria consisted of cases available with complete data, studies published only in English, studies that dealt with adults who were diabetic with essential hypertension and the

studies that were cross-sectional in nature and those that were reviews and meta analyses conducted in Asia.

The exclusion criteria consisted of case reports, studies that included Non-Asian countries, paediatric and pregnant population, studies that included patients with secondary hypertension and those that were published in languages other than English. Studies that included insulin dependent diabetic patients were not taken and those that had a sample size less than 200 were also not considered.

No ethical approval was needed since we conducted a systematic review and meta-analyses.




Figure 1: Flowchart for inclusion of studies in the systematic review and meta analyses

Statistical Analysis

Statistical analysis was performed with the SPSS version 28.0 and the data were presented using descriptive statistics such as number and percentage. R Studio was used for the preparation of graphs.

Results

Screening Flow

Usable data from eligible studies were independently extracted by two authors (S.S and A.T.C.). Any discrepancies in their decisions were resolved through consultation with a third author (J.K.S.). During the duplicate removal stage, 590 articles were removed. Out of a total of 26613 articles, 25600 articles were excluded during title and abstract screening. Finally, a total of 12 articles were considered for the systematic review and meta-analyses after excluding 1001 articles from the remaining 1013 articles during full text screening process.

Study Characteristics

This systematic review and meta-analyses included a total of 12 studies.

The studies were analysed by using Microsoft Excel 2016 and R Studio was used for graphical preparations. The two authors

(S.S and A.T.C.) assessed the methodology and quality by using the New Castle Ottawa Scale (Wells GA *et al*, 2000). Finally, a total of 12 studies met the quality of assessment. Our review included studies from 12 Asian countries namely India, Thailand, Kuwait, Lebanon, Egypt, Sri Lanka, Iran, Bangladesh, Jordan, China, Pakistan and Saudi. The author names, country, type of study, period of study, sample size and prevalence of hypertension in diabetic patients were tabulated (**Table 1 and Figure 2**). The studies selected were from various countries of Asia to evaluate the prevalence of hypertension in the diabetic population. Forest plot was prepared for prevalence of hypertension in diabetics in Asia stated by various authors (**Figure 2**). The pooled prevalence was 0.60(95 CI – 0.49 to 0.70). In this study, the heterogeneity was tested with $I^2 = 99.841\%$ and P value <0.001 indicating the presence of heterogeneity.

Funnel test and Egger's test

Funnel plot was found out to be roughly symmetrical indicating minor publication bias (**Figure 3**). Egger's test was also used to rule out the bias (Egger M *et al*, 1997). The results showed that P value is 0.341 that is >0.05 suggesting that there was publication bias.

Kuwait and China reported relatively high prevalence of hypertension in diabetic patients followed by Sri Lanka, Lebanon and Jordan. Saudi reported the lowest hypertension prevalence in diabetic patients followed by Iran (**Table 1 and Figure 4**).

Table 1: Study characteristics and HTN prevalence in T2DM patients across countries in Asia

SI No	Author	Country	Type of Study	Period of Study	Prevalence (%)	Sample Size
1	Dhananjay Yadav <i>et al</i> , I 2014	India	Cross-sectional	3 years	49%	700
2	Cameron Hurst <i>et al</i> . 2015	Thailand	Cross-sectional	3 years	55.35%	55797
3	Arshad Mohamed Channanath <i>et al</i> , I 2015	Kuwait	Cross-sectional, retrospective	12 years	87.16%	3904
4	Lama Soubra <i>et al</i> , I 2018	Lebanon	Cross-sectional, retrospective	1 year	75%	700
5	Eman H Waly <i>et al</i> , I 2018	Egypt	Cross-sectional	3 months	68%	300
6	Maulee Hiromi Arambewela <i>et al</i> , I 2018	Sri Lanka	Cross-sectional	7 months	77.6%	3000
7	Soghra Rabizadeh <i>et al</i> , I 2021	Iran	Cross-sectional	1 year	36.1%	2612
8	Hiba Alsaadon <i>et al</i> , I 2022	Bangladesh	Cross-sectional, retrospective	1 year	67.2%	1252
9	Ahlam Bani Salameh <i>et al</i> , I 2022	Jordan	Cross-sectional	2 years	74.6%	1382
10	Xin Zhang <i>et al</i> , I 2024	China	Systematic review and meta-analyses	19 years	87.16%	7517
11	Mehwish Javeed <i>et al</i> , I 2024	Pakistan	Review	--	47.06%	2327
12	Abdulhameed A Alharbi <i>et al</i> , I 2024	Saudi	Retrospective. Cohort	5 years	23%	274

Table 2: Triggering factors

SI No	Authors	Others
1	Dhananjay Yadav <i>et al</i> , 2014	Dyslipidemia Males- 56.3%, Females- 72% Metabolic syndrome 6-10 years 39.4% SBP: 60.5% DBP: 80%
2	Arshad Mohamed Channanath <i>et al</i> , I 2015	Obesity 29.6%
3	Lama Soubra <i>et al</i> , I 2018	Dyslipidemia 75.6%
4	Eman H Waly <i>et al</i> , I 2018	Physical inactivity 97% Obesity 76.4% Cholesterol 62.1% Family history 65%
5	Maulee Hiromi Arambewela <i>et al</i> , I 2018	Vascular complication 84.4% Dyslipidemia 76.7% Obesity 75.7%
6	Soghra Rabizadeh <i>et al</i> , I 2021	Obesity 39.3% Metabolic Syndrome 88.8%
7	Hiba Alsaadon <i>et al</i> , I 2022	Physically inactive 69.6% Obesity 75.1% High waist hip ratio 63.3%
8	Ahlam Bani Salameh <i>et al</i> , I 2022	Hip waist circumference ratio 89.9%

*SBP: Systolic Blood Pressure; DBP: Diastolic Blood Pressure

Table 3: Correlation of hypertensive diabetes with gender

SI No	Authors	Gender	
		Male	Female
1	Dhananjay Yadav <i>et al</i> , I 2014	55.2%	42.9%
2	Arshad Mohamed Channanath <i>et al</i> , I 2015	41%	59%
3	Lama Soubra <i>et al</i> , I 2018	55.2%	44.8%
4	Eman H Waly <i>et al</i> , I 2018	42.4%	57.6%
5	Soghra Rabizadeh <i>et al</i> , I 2021	34.3%	65.7%
6	Hiba Alsaadon <i>et al</i> , I 2022	63.1%	64.9%
7	Ahlam Bani Salameh <i>et al</i> , I 2022	48.05%	51.95%
8	Xin Zhang <i>et al</i> , I 2024	59.8%	40.2%

Table 4: Correlation of hypertensive diabetes with urbanization

SI No	Authors	Urbanization	
		Urban	Rural
1	Hiba Alsaadon <i>et al</i> , I 2022	68.2%	54.2%
2	Ahlam Bani Salameh <i>et al</i> , I 2022	63.65%	36.35%

Table 5: T2DM prevalence in smokers

SI No	Authors	Smoking
1	Lama Soubra <i>et al</i> , I 2018	56%
2	Maulee Hiromi Arambewela <i>et al</i> , I 2018	11%
3	Soghra Rabizadeh <i>et al</i> , I 2021	10.30%
4	Hiba Alsaadon <i>et al</i> , I 2022	66.10%

Table 6: Merits and gaps

SI No	Authors	Merits	Gaps
1	Dhananjay Yadav <i>et al</i>	Robust dataset	Short period of study (3 years)
2	Cameron Hurst <i>et al</i>	First nationwide multicentre study in South East Asia with large sample	Absence of lifestyle and dietary data may have led to potential information bias.
3	Arshad Mohamed Channanath <i>et al</i>	First report to assess age and BMI correlation in T2DM hypertensive patients	Family history, lifestyle factors and dietary patterns were missing contributing to significant bias
4	Lama Soubra <i>et al</i>	one of the few studies that addressed hypertension in diabetes in Middle East	BP was recorded once adding to potential bias. Factors contributing to low controlled BP levels like medications and less stringent goals were not included.
5	Eman H Waly <i>et al</i>	Robust dataset	Cross-sectional nature leading to non-establishment of cause-effect relationship.
6	Maulee Hiromi Arambewela <i>et al</i>	Large scale descriptive study conducted in National Hospital of Sri Lanka which is the largest tertiary care hospital in the country allowing diverse representative sample of 3000 contributing to the reliability of the findings	Prevalence might have been over reported.
7	Soghra Rabizadeh <i>et al</i>	multivariate logistic regression modelling was used to recognize the independent factors of uncontrolled hypertension that strengthened the findings of the study. metabolic syndrome impacting hypertension in diabetic patients was depicted.	Cross-sectional nature
8	Hiba Alasaadon <i>et al</i>	Robust sample data from over 6 diabetes hospitals providing primary, secondary and tertiary care covering urban and rural settings utilized.	Cross-sectional nature
9	Ahlam Bani Salameh <i>et al</i>	Use of standardized automated sphygmomanometers ensured accurate data collection. Sociodemographic factors, anthropometric measurements, HbA1c levels and lipid profiles, contributed to thorough analysis of factors associated with HTN	Data missed out important factors such as medication adherence and patient behaviour.
10	Xin Zhang <i>et al</i>	Reported as first systematic review to investigate hypertension in T2DM Chinese patients.	High heterogeneity was noted and small number of studies were included.

11	Mehwish Javeed et al	Risk factors such as dyslipidaemia, obesity and poor lifestyle choices, high sodium intake and sedentary life were identified.	Generalizability limited due to varying health outcomes on basis of demographics and regions.
12	Abdulhameed A Alharbi et al	Thorough data collection was done using customized checklists using various demographic factors and health indicators.	Retrospective might have relied on existing health records only that might not always be very accurate.




Figure 2: Forest plot for prevalence of HTN in diabetic Asians author wise




Figure 3. Funnel plot for HTN prevalence in Diabetic Asians




Figure 4: Choropleth map depicting HTN prevalence in Diabetic Asians




Figure 5: Stacked area chart depicting various triggering factors of diabetes author wise




Figure 6: Dumbbell plot depicting male vs female HTN prevalence in diabetic Asians




Figure 7: Association of urbanization and T2DM




Figure 8: Bubble heat map representing smoking and T2DM association author wise



Figure 9: Inter-relation between T2DM, HTN and various triggering factors causing morbidity

Discussion

A critical area of research is the interrelationship between type 2 diabetes mellitus and hypertension, and this is evident in the findings of different authors of our study.

The correlation between hypertensive diabetes and various triggering factors was tabulated (**Table 2, 3, 4 & 5; Figure 5,6,7 & 8**).

The significant role of dyslipidaemia and metabolic syndrome in the prevalence of hypertension among diabetic patients was underlined by an author pointing out that these factors are the causes of the increased cardiovascular risk associated with diabetes (Dhananjay Yadav *et al*, 2014). This was supported by another study (Suanrueang P, 2024). Another author of our study who further reported that hypertension is associated with microvascular complications in diabetic patients, found the notion that the presence of hypertension further complicates such problems as retinopathy and nephropathy (Cameron Hurst *et al*, 2015). This was further corroborated on by another author (Joshi BS *et al*, 2024). Another author of our study added to this argument by identifying obesity as the key factor leading to hypertension in diabetic patients and hence stated that the increasing rates of obesity in the region are directly linked to the simultaneous increase in the incidence of both conditions (Arshad Mohamed Channanath *et al*, 2015). This was further elucidated upon by another author (Pasdar Y *et al*, 2024). According to an author in our study the impact of physical inactivity, dyslipidaemia and smoking are the major contributors to hypertension in Lebanon, which supports the notion that lifestyle factors are the key to the management of both diseases (Lama Soubra *et al*, 2016). This was further showed in another study (Yang X *et al*, 2024). Another author of our study agreed with these findings, saying that in addition to physical inactivity, obesity is one of the major risk factors in Egypt, all of which complicate the proper management of hypertension among diabetic patients (Eman Waly *et al*, 2018).

Yet another author of our study mentioned the prevalence of the alarming vascular complications among diabetic patients with hypertension in Sri Lanka, and stressed upon the fact that there was a link of acute health consequences with the two conditions dropping down the patients to irreversible stage (Maulee Hiromi Arambewela *et al*, 2018). One more author of our study created awareness of the importance of metabolic syndrome which is a common factor in diabetic patients who had hypertension in Iran (Soghra Rabizadeh *et al*, 2021). This was depicted in another study (Vareldzis R *et al*, 2024). Two authors of our study stressed on urbanization, lifestyle changes and hip waist circumference ratio in Bangladesh by depicting their impact on increasing hypertension among diabetic patients (Hiba Alsaadon *et al*, 2022; Ahlam Bani Salameh *et al*, 2022). This was further noted in another study (Talukder A *et al*, 2024). Yet another author demonstrated hyperinsulinemia as a biochemical pathway for the cause of hypertension in diabetic people (Xin Zhang *et al*, 2024). This was reported by another author (Fazio S *et al*, 2024). Yet another author of our study emphasized on the poor lifestyle choices and dyslipidemia as the important factors existing in Pakistan stressing on the need to adopt comprehensive lifestyle interventions to tackle both diabetes and high blood pressure (Mehwish Javeed *et al*, 2024). Finally, another author of our study observed the regional differences in the prevalence of hypertension among diabetic patients in Saudi Arabia by recommending the authorities to respond with proper localized strategies to control and eliminate these interconnected conditions (Abdulhameed A Alharbi *et al*, 2024). Together, these researches reveal the multifactorial nature regarding the relationship between

T2DM and HTN by supporting the fact that lifestyle adjustments, metabolic syndrome and urbanization are the triggering factors. The merits and gaps of each study by various authors of our study were tabulated (**Table 6**).

Conclusion

The drivers of hypertension in a diabetic patient leads to dreadful catastrophe, hence awareness should be created among the common man. There should be facilities available to the public for cheap availability of digital BP monitor supply through ration shops. No smoking sign boards must be placed in areas in malls and public areas. All means to increase the awareness of this deadly duo namely HTN and T2DM to the public is made available through social media, advertisements on TV, and movies. NGO, and other organisations can arrange free camps and health check -up for the public. Our study aimed to evaluate the relationship between T2DM and HTN across various countries in Asia and showed positive correlation between the two conditions. The various triggering factors like obesity, dyslipidaemia, sedentary lifestyle, smoking and high sodium intake are the tip of the iceberg (**Figure 9**).

Declarations

Ethical Approval

Not Required

Source of Funding

This research was not supported by any specific grants from public, commercial, or non-profit funding agencies.

Data Availability

Available upon responsible request by corresponding author.

Conflicts of Interests

The authors report no conflict of interest.

Author Contributions

Conceptualization and methodology, S.S., A.T.C, and J.K.S; Formal analysis, A.T.C, S.S., J.K.S.; Visualization and writing – original draft S.S., A.T.C., J.K.S., H.A.; Writing – review and editing, S.S., A.T.C., J.K.S., H.A. and J.H. All authors have read and agreed to the final version of the manuscript.

Acknowledgments

We would like to thank our Principal, Dr. Vasanthamalai, and General Manager, Mr. Rahim for their immense involvement. And Miss. Swathi for her technical assistance with data analysis and illustrations in the preparation of this study.

References

- [1] Zimmet P. The burden of type 2 diabetes: are we doing enough?. *Diabetes & metabolism*. 2003 Sep 1;29(4):6S9-18.
- [2] Wild S, Roglic G, Green A, Sicree R, King H: Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*, 27:1047–1053, 2004

- [3] Wang Z, do Carmo JM, da Silva AA, Fu Y, Hall JE. Mechanisms of synergistic interactions of diabetes and hypertension in chronic kidney disease: role of mitochondrial dysfunction and ER stress. *Current hypertension reports*. 2020 Feb;22:1-0.
- [4] Cheung BM. The hypertension–diabetes continuum. *Journal of cardiovascular pharmacology*. 2010 Apr 1;55(4):333-9.
- [5] Parving HH, Hommel E, Mathiesen E, Skøtt P, Edsberg B, Bahnson M, Lauritzen M, Hougaard P, Lauritzen E. Prevalence of microalbuminuria, arterial hypertension, retinopathy, and neuropathy in patients with insulin dependent diabetes. *Br Med J (Clin Res Ed)*. 1988 Jan 16;296(6616):156-60.
- [6] Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, Stein C, Basit A, Chan JC, Mbanya JC, Pavkov ME. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes research and clinical practice*. 2022 Jan 1;183:109119.
- [7] Cohen B. Urbanization in developing countries: Current trends, future projections, and key challenges for sustainability. *Technology in society*. 2006 Jan 1;28(1-2):63-80.
- [8] Devi BN, Kumar MV, Sreedhar M. Prevalence of risk factors for Non Communicable Diseases in urban slums of Hyderabad, Telangana. *Indian Journal of Basic and Applied Medical Research*. 2014;4(1):487-93.
- [9] Sechi LA, Bartoli E. Mechanisms of insulin resistance leading to hypertension: what we can learn from experimental models. *J Investig Med*. 1997 Jun;45(5):238-51. PMID: 9249996.
- [10] Moher D, Liberati A, Tetzlaff J, Altman DG. Research methods and reporting. *Bmj*. 2009 Aug 8;8:332-6.(PRISMA)
- [11] Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses. 2000
- [12] Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *bmj*. 1997 Sep 13;315(7109):629-34.
- [13] Yadav D, Mishra M, Tiwari A, Bisen PS, Goswamy HM, Prasad GB. Prevalence of dyslipidemia and hypertension in Indian type 2 diabetic patients with metabolic syndrome and its clinical significance. *Osong Public Health Res Perspect*. 2014 Jun;5(3):169-75.
- [14] Suanrueang P. A comparison of the disease occurrence of cerebrovascular diseases, diabetes mellitus, hypertensive diseases, and ischaemic heart diseases among hospitalized older adults in Thailand. *Scientific Reports*. 2024 Jan 2;14(1):123.
- [15] Hurst C, Thinkhamrop B, Tran HT. The Association between Hypertension Comorbidity and Microvascular Complications in Type 2 Diabetes Patients: A Nationwide Cross-Sectional Study in Thailand. *Diabetes Metab J*. 2015 Oct;39(5):395-404.
- [16] Joshi BS, Paranjpe GS. Connection Between Type 2 Diabetes and Diabetic Retinopathy, Hypertension, Dyslipidemia, and Nephropathy. In *Advancements in Science and Technology for Healthcare, Agriculture, and Environmental Sustainability* 2024 Jun 13 (pp. 410-416). CRC Press.
- [17] Channanath AM, Farran B, Behbehani K, Thanaraj TA. Association between body mass index and onset of hypertension in men and women with and without diabetes: a cross-sectional study using national health data from the State of Kuwait in the Arabian Peninsula. *BMJ Open*. 2015 Jun 4;5(6):e007043.
- [18] Pasdar Y, Rezaeian S, Mohammadi E, Khosravi shadmani F, Shahnazi N, Najafi F, Nazar MM, Darbandi M. The interaction between general or abdominal obesity and hypertension on the risk of type 2 diabetes mellitus: a cross-sectional analysis in Iranian adults from the RaNCD cohort study. *BMC Public Health*. 2024 Mar 11;24(1):752.
- [19] Soubra LA, Nureddin HA, Omar AG, Saleh MO. Factors associated with hypertension prevalence and control among Lebanese type 2 diabetic patients. *Int J Pharm Pharm Sci*. 2016;8(10):153-9.
- [20] Yang X, Sun J, Zhang W. Global trends in burden of type 2 diabetes attributable to physical inactivity across 204 countries and territories, 1990-2019. *Frontiers in Endocrinology*. 2024 Feb 26;15:1343002.
- [21] Waly EH, Hamed MS. Hypertension and dyslipidemia among type ii diabetic patients and related risk factors and complications. *Egypt J Commun Med*. 2018 Jan;36(1):31-43.
- [22] Arambewela MH, Somasundaram NP, Jayasekara HBPR, Kumbukage MP, Jayasena PMS, Chandrasekara CMPH, Fernando KRAS, Kusumsiri DP. Prevalence of Chronic Complications, Their Risk Factors, and the Cardiovascular Risk Factors among Patients with Type 2 Diabetes Attending the Diabetic Clinic at a Tertiary Care Hospital in Sri Lanka. *J Diabetes Res*. 2018 May 23;2018:4504287. doi: 10.1155/2018/4504287. PMID: 29951551; PMCID: PMC5989171.
- [23] Rabizadeh S, Gholami B, Mahmoudzadeh Kani S, Rajab A, Farrokhpour H, Esteghamati A, Nakhjavani M. Uncontrolled hypertension in patients with type 2 diabetes: What are the correlates?. *The Journal of Clinical Hypertension*. 2021 Sep;23(9):1776-85.
- [24] Varelzdis R, Perez A, Reisin E. Hyperuricemia: an intriguing connection to metabolic syndrome, diabetes, kidney disease, and hypertension. *Current Hypertension Reports*. 2024 Jan 25:1-9.
- [25] Alsaadon H, Afroz A, Karim A, Habib SH, Alramadan MJ, Billah B, Shetty AN. Hypertension and its related factors among patients with type 2 diabetes mellitus - a multi-hospital study in Bangladesh. *BMC Public Health*. 2022 Jan 29;22(1):198.
- [26] Salameh AB, Hyassat D, Suhail A, Makahleh Z, Khader Y, El-Khateeb M, Ajlouni K. The prevalence of hypertension and its progression among patients with type 2 diabetes in Jordan. *Ann Med Surg (Lond)*. 2021 Dec 8;73:103162.
- [27] Talukder A, Sara SS, Hossain MT, Nath CD, Rahman R, Hussain S, Sarma H, Huda MN. Rural and urban differences in the prevalence and determinants of Type-2 diabetes in Bangladesh. *Plos one*. 2024 Apr 11;19(4):e0298071.
- [28] Zhang X, Yang XL, Liu S, Ye YS, Yang QT, Xia LN. Prevalence of hypertension among type 2 diabetes mellitus patients in China: a systematic review and meta-analysis. *Int Health*. 2024 Mar 4;16(2):144-151.

- [29] Fazio S, Affuso F, Cesaro A, Tibullo L, Fazio V, Calabrò P. Insulin Resistance/Hyperinsulinemia as an Independent Risk Factor That Has Been Overlooked for Too Long. *Biomedicines*. 2024 Jun 26;12(7):1417.
- [30] Javed M, Asim M, Ishfaq H, Ikram N, Shamim U, Shahid M, Syed SK. Prevalence of hypertension in diabetic patients in Pakistan. *Journal of Contemporary Pharmacy*. 2024 Jun 30;8(1):28-32.
- [31] Alharbi AA, Alharbi AA, Al-Dubai SA. Inter-relation between diabetes mellitus and hypertension in terms of incidence and prediction in Saudi Arabia: a retrospective cohort study. *BMC Public Health*. 2024 Jul 22;24(1):1956.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025